



## Standard Operating Procedure: Mouse Oral Gavage

These SOPs were developed by the Office of the University Veterinarian and reviewed by Virginia Tech IACUC to provide a reference and guidance to investigators during protocol preparation and IACUC reviewers during protocol review. They can be used as referenced descriptions for procedures on IACUC protocols.

However, it is the sole responsibility of the Principal Investigator to ensure that the referenced SOPs adequately cover and accurately represent procedures to be undertaken in any research project. Any modification to procedure as described in the SOP must be outlined in each IACUC protocol application (e.g. if the Principal Investigator plans to use a needle size that is not referenced in the SOP, simply state that alteration in the IACUC protocol itself).

### Table of Contents

- I. Procedure Summary and Goal ..... 2
- II. Personal Protective Equipment (PPE) and Hygiene ..... 2
- III. Supply List ..... 2
- IV. Detailed Procedure ..... 2
- V. Variations ..... 4
- VI. Potential Adverse Effects, Mitigation, or Treatment ..... 4
- VII. References ..... 4

## I. Procedure Summary and Goal

Describes the procedure for the administration of fluids directly into the lower esophagus or stomach through the use of a standard gavage needle (or feeding needle) introduced into the mouth and threaded down the esophagus.

Considerations:

- a. With proper training, this method is performed easily and quickly with minimal distress to the animal. Firm manual restraint with immobilization of the head is necessary.
- b. The position of the mouse is very important. Align the head and body vertically with the esophagus.
- c. Never force the feeding needle down, allow the mouse to swallow and gravity to assist.
- d. Always administer the compound slowly and finish administering before pulling gavage needle out. Always pull needle straight out. Do Not Aspirate!
- e. STOP the procedure if mucous membranes are blue, the mouse struggles vigorously, or fluid comes from the nose
- f. Precoating the gavage needle with sucrose has been demonstrated to decrease stress-related reactions and thereby improve animal welfare during oral gavage. (J Am Assoc Lab Anim Sci. 2010 May;49(3):329-34)

## II. Personal Protective Equipment (PPE) and Hygiene

- a. Ensure appropriate PPE is used to protect technician from accidental exposure to blood and other body fluids, such as:
  - i. Gloves
  - ii. Eye protection
  - iii. Mask
  - iv. Other PPE as required by protocol/facility
- b. Hands should be washed and/or gloves changed between animals.
- c. Promptly dispose of used sharps in the provided leak-proof, puncture resistant sharps container.

## III. Supply List

- a. Prefilled syringes
- b. Gavage needle (Figure 1)
  - i. Reusable: stainless steel (ball tip, shaft, and hub)
  - ii. Single-use: silicone ball tip, stainless steel shaft, plastic hub

## IV. Detailed Procedure

- a. Frequency and Volume

- i. Up to 10 ml/kg
- ii. Dosing can be repeated up to three times in a 24-hour period
  - 1. If additional dosing is necessary, it must be justified in the animal use protocol.
- b. Anesthesia
  - i. General anesthesia is not recommended and may interfere with gastric emptying.
- c. Procedure
  - i. Select appropriate size gavage needle based upon weight and size of animal.

Weight (gms)	Gauge	Length (inch)	Ball diameter (mm)	Shape
to 14	24	1"	1.25	Straight
15-20	22	1-1.5"	1.25	Straight
20-25	20	1-3	1.9-2.25	Straight or Curved
25-30	18	1-2	2.0-2.3	Straight or Curved
30-35	18	2-3	2.0-2.3	Straight or Curved
Suggested needle sizes for mice (as listed in Braintree Scientific Catalog)				

- ii. Place the gavage needle along the lateral aspect of the animal so that the ball tip of the gavage needle is adjacent to the last rib. Note where the shaft of the needle lines up with incisors to determine how far needle can be safely advanced (Figure 2).
- iii. To restrain, securely scruff the animal and hold its head up with the longitudinal axis of the body in a vertical position; animal should face handler with head immobilized (Figure 3). This position allows gravity to assist in preventing reflux of the material being administered.



Figure 2. Measuring a Gavage Needle



Figure 3. Restraint Presentation




Figure 4. Introducing the Gavage Needle Tilting Head Back



Figure 5. Gavage Needle Fully Inserted

- iv. Insert the gavage needle into the mouth and direct over the tongue and into the pharynx. Slightly hyperextending the animal's head back will facilitate entry of the gavage needle into the esophagus (Figure 4).
  1. The animal will swallow by reflex when the pharynx is entered and the needle should easily slide into the esophagus with minimum pressure. Do not force needle.
  2. The ball tip should prevent gavage needle from entering the trachea, but if animal gasps, remove needle into oral cavity and repeat procedure of passing the needle over the tongue.
  3. Note depth of the gavage needle at the appropriate site/mark from previous measurement.
- v. If animal breathes without struggling, administer amount of material slowly and smoothly (Figure 5).
- vi. Once administration completed, pull gavage needle straight out.
- vii. Observe animal for normal respiration.
  1. Observe for signs of labored breathing or distress
  2. If the animal gasps for air, immediately euthanize.

## V. Variations

None

## VI. Potential Adverse Effects, Mitigation, or Treatment

- a. Tracheal administration
  - i. Stop administration of substance, remove gavage needle
  - ii. Contact veterinary personnel immediately
- b. Reflux, aspiration and/or respiratory distress
  - i. Stop administration of substance, remove gavage needle
  - ii. Contact veterinary personnel immediately
- c. Pharyngeal, esophageal, gastric irritation, injury or rupture
  - i. Contact veterinary personnel immediately
- d. Death

## VII. References

American Association of Laboratory Animal Science. Laboratory Animal Technician Training Manual. (Memphis, TN: Drumwright and Co, 2007)

Charles River Insourcing Solutions. Biomethodology in the Laboratory Mouse

Charles River SOP 2405-3 – Dosing of Rodents – TGS and Discovery Services

Hedrich, H., Bullock, G., and Petrusz, P., *The Laboratory Mouse*, (London, England: Elsevier Academic Press, 2004)

Hoggatt, A.F., Hoggatt, J., Honerlaw, M., and Pelus, L.M. A Spoonful of Sugar Helps the Medicine Go Down: A Novel Technique to Improve Oral Gavage in Mice. *J Am Assoc Lab Anim Sci.* 49(3), 329-34 (May 2010).

National Research Council. *Guide for the Care and Use of Laboratory Animals*. (Washington, D.C.: National Academy Press, 2011)

Suckow, M., Danneman, P., and Brayton, C. *The Laboratory Mouse*. (Boca Raton, FL: CRC Press LLC, 2001)

Turner, P.V., Brabb, T., Pekow, C., and Vasbinder, M. Administration of Substances to Laboratory Animals: Routes of Administration and Factors to Consider. *J Am Assoc Lab Anim Sci.*; 50(5): 600–613. (2011 September) <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3189662/>